



# Postnatal depression

## A guide for women and their families

After the birth of a baby women experience many physical, emotional and social changes. This makes the chance of experiencing depression higher. So, early parenthood is a time when it is important to watch for its signs and symptoms. If you think you may have depression, seek help early so you can take steps to get on top of the depression.

### How common is postnatal depression?

Up to **1 in 7** Australian women feel depressed in the first year after having a baby. It is common to experience depression and anxiety at the same time.

### What causes postnatal depression?

There is no single cause of depression. It is likely to be a result of:

- your genetic make-up and family history
- your personal way of thinking and coping
- features of your environment that influence your mental and emotional wellbeing.

A combination of these factors may make your risk of developing depression higher. There are also factors that protect you, such as having good support networks.

### What are the signs and symptoms?

Many women miss the early signs of depression - sometimes because they think it is the 'baby blues' or adjusting to parenting (e.g. tiredness). It is important to be aware of common symptoms including:

- feeling low or numb - some women describe feeling nothing at all
- loss of confidence, feeling helpless, hopeless and worthless
- feeling teary and emotional, angry, irritable or resentful towards others
- changes in sleep - not being able to sleep even when you have the chance or wanting to sleep all the time
- changes in appetite - accompanied by weight loss or weight gain
- lack of interest and/or energy
- difficulties concentrating, thinking clearly or making decisions (which can also result from lack of sleep)
- feeling isolated, alone and disconnected from others
- having thoughts of harming yourself, your baby and/or other children
- finding it difficult to cope and get through the day.

### How is postnatal depression identified and diagnosed?

Your doctor or maternal and child health nurse may offer you screening with the Edinburgh Postnatal Depression Scale and/or the Antenatal Risk Questionnaire.

- The Edinburgh Postnatal Depression Scale is series of 10 questions about how you have been feeling in the last 7 days. For each question, you choose the response that describes your feeling.
- The Antenatal Risk Questionnaire asks questions about your life more generally and helps to identify factors that may place you at greater risk of experiencing depression.

If you think you might be experiencing depression, you can ask your doctor or maternal and child health nurse for these screening assessments. Your score(s) will tell you and your doctor or maternal and child health nurse whether it would be helpful to further assess your mental health.

A GP or trained mental health professional can diagnose depression. They do this by asking whether you have had certain symptoms over a period of time (usually 2 weeks or more).

### How is postnatal depression treated?

Your doctor will talk to you about psychological (talking) therapies and possibly medicines for treating depression. You will also benefit from emotional and physical support. The type of treatment that is right for you will depend on your situation, your symptoms and the treatments you can access. The sooner you seek help, the sooner you can work towards recovery. Getting help early is best for both you and your baby.

#### Talking therapies

Talking therapies can be very helpful if treating symptoms of depression. Talking therapies can help you recognise and change the negative thinking and feelings that depression brings. They also give you useful tools to stop depression coming back.

# Postnatal depression

## A guide for women and their families



Talking therapies include:

- Cognitive-behavioural therapy (CBT): the cognitive (or thinking) part of this therapy teaches you to think logically and challenge negative thoughts. The behaviour part helps you change the way you react in situations and can help you to get involved in activities that you have been avoiding or have stopped doing
- Interpersonal therapy (IPT): helps you to find new ways to connect with others, and overcome losses, challenges and conflicts that you may have.

Registered practitioners with appropriate training and experience provide talking therapies for individuals or groups.

### Medicines

Depending on how helpful you find talking therapies, you may also need to take medicine. Talking therapies may continue to be helpful as your symptoms begin to improve.

Medicines for depression are called antidepressants. There are different types of medicine and they work in different ways. Selective serotonin reuptake inhibitors (SSRIs) can be safely used when you are breastfeeding and can also help with anxiety symptoms. Your doctor may also suggest another type of medicine called tricyclic antidepressants, especially if they have worked for you in the past.

Some useful tips about taking antidepressants.

- The medicine can take up to 3 weeks to have an effect so other supports can be helpful in the meantime.
- You may experience some mild side-effects - talk to your doctor if these become too hard to cope with.
- You may need to try more than one medicine to find the one that is best for you.
- Most women stay on their medicine for 6-24 months - don't stop taking a medicine suddenly once you start feeling better.
- When you come off the medicine, you should do this slowly - your doctor will help you and check for any side-effects or symptoms of relapse.

### Electroconvulsive therapy (ECT)

ECT is used when symptoms of depression are severe, medicine is not working and the risks of symptoms (e.g. thoughts of self-harm) are very high.

## Advice for women experiencing postnatal depression

### Seek help and treatment

- Seek help from a doctor or other health professional.
- Learn about effective treatments.
- Call a support service or mental health crisis line if other help is not available.

### Accept help and support

- Develop a support system of friends, family and professionals and accept help.
- Discuss your feelings with your partner, family and friends.

### Look after your physical health

- Try to eat healthy meals, including fruit, vegetables, whole grains and lots of water.
- Plan some enjoyable physical activity every day.
- Try to establish good sleeping patterns.
- Practice techniques to reduce stress, such as muscle relaxation and deep breathing.

## Advice for family and friends providing support

### Listen and reassure

- Encourage the woman to discuss her symptoms.
- Explain that depression can be treated and managed.

### Provide information

- Give the woman good quality information about postnatal depression, such as this fact sheet.
- Give details of helplines if she is feeling distressed and needs support.
- Offer information to the woman's partner/others.



# Postnatal depression

## A guide for women and their families

### Direct to care and support

- Encourage the woman to talk to her GP or other health professional.
- Encourage the woman to identify and draw on possible supports and services that may be available to her for practical and/or emotional support.
- Remind the woman that she can go to her doctor or local hospital if she is at risk of harming herself or others.

### Information for women and their families:

#### Ready to COPE Guide:

Sign up to receive free weekly information about emotional and mental health throughout your pregnancy and first year with a new baby, at [readytcope.org.au](https://www.readytcope.org.au)

#### To talk to someone:

To speak with a specialist counsellor, call the [PANDA helpline](https://www.panda.org.au) on 1300 726 306 (Monday to Saturday 9.00am – 7.30pm AEST/AEDT)

#### To find a service:

To find out about other perinatal mental health treatment and support services, visit the [eCOPE Directory](https://www.eCOPE.org.au)

### Where you can get more information:

More information about depression in early parenthood can be found at [cope.org.au](https://www.cope.org.au). This includes information on the following topics:

[What increases my risk?](#)

[About postnatal depression](#)

[Treatment of postnatal depression](#)

[When to seek help](#)

[Available support under Medicare](#)

[Mental health under Medicare](#)

[Helpline, Resources and Services](#)

[Find help in my area for postnatal depression](#)



**COPE**  
Centre of Perinatal Excellence

This fact sheet has been developed by COPE: Centre of Perinatal Excellence and is derived from *Mental Health Care in the Perinatal Period: Australian Clinical Practice Guideline*. 2023. Centre of Perinatal Excellence (COPE).

Funded by the Australian Government Department of Health and Aged Care.